



## SWFSC “SOIL CRITTERS” FACT SHEET

### REFERENCES

- A’Bear, A. D., Boddy, L., Raspopnig, G., & Jones, T. H. (2010). Non-trophic effects of oribatid mites on cord-forming basidiomycetes in soil microcosms. *Ecological Entomology*, 35(5), 483–490. <https://doi.org/10.1111/j.1365-2311.2010.01204.x>
- Anderson, J. M. (1975). The enigma of soil animal species diversity. In J. Vaněk (Ed.), *Progress in Soil Zoology* (pp. 51–58). Springer Netherlands.
- Griffiths, B. S. (1994). Microbial-feeding nematodes and protozoa in soil: Their effects on microbial activity and nitrogen mineralization in decomposition hotspots and the rhizosphere. *Plant and Soil*, 164(1), 25–33.
- Keller, T., & Or, D. (2022). Farm vehicles approaching weights of sauropods exceed safe mechanical limits for soil functioning. *Proceedings of the National Academy of Sciences of the United States of America*, 119(21), e2117699119.
- Klironomos, J. N., & Hart, M. M. (2001). Animal nitrogen swap for plant carbon. *Nature*, 410(6829), 651–652.
- Maraun, M., Visser, S., & Scheu, S. (1998). Oribatid mites enhance the recovery of the microbial community after a strong disturbance. *Applied Soil Ecology: A Section of Agriculture, Ecosystems & Environment*, 9(1–3), 175–181.
- Menzel, R., Geweiler, D., Sass, A., Simsek, D., & Ruess, L. (2018). Nematodes as important source for omega-3 long-chain fatty acids in the soil food web and the impact in nutrition for higher trophic levels. *Frontiers in Ecology and Evolution*, 6, 96. <https://doi.org/10.3389/fevo.2018.00096>
- Neher, D. A., Weicht, T. R., & Barbercheck, M. E. (2012). Linking invertebrate communities to decomposition rate and nitrogen availability in pine forest soils. *Applied Soil Ecology: A Section of Agriculture, Ecosystems & Environment*, 54, 14–23.
- Neher, D. A. (2010). Ecology of plant and free-living nematodes in natural and agricultural soil. *Annual Review of Phytopathology*, 48(1), 371–394.
- Norton, R. A., & Behan-Pelletier, V. M. (1991). Calcium carbonate and calcium oxalate as cuticular hardening agents in oribatid mites (Acari: Oribatida). *Canadian Journal of Zoology*, 69(6), 1504–1511.

Orgiazzi, A., Bardgett, R. D., Barrios, E., Behan-Pelletier, V., Briones, M. J. I., De Deyn, G. B., Eggleton, P., Fierer, N., Fraser, T., Hedlund, K., Jeffrey, S., Johnson, N. C., Jones, A., Kandeler, E., Kaneko, N., Lavelle, P., Lemanceau, P., Miko, L., Montanarella, L., ... Wall, D. H. (2016). *Global Soil Biodiversity Atlas*. European Commission, Publications Office of the European Union.

Ortega-Jimenez, V. M., Challita, E. J., Kim, B., Ko, H., Gwon, M., Koh, J.-S., & Bhamla, M. S. (2022). Directional takeoff, aerial righting, and adhesion landing of semiaquatic springtails. *Proceedings of the National Academy of Sciences of the United States of America*, 119(46), e2211283119.

Pingree, M. R. A., & Kobziar, L. N. (2019). The myth of the biological threshold: A review of biological responses to soil heating associated with wildland fire. *Forest Ecology and Management*, 432, 1022–1029. <https://doi.org/10.1016/j.foreco.2018.10.032>

Platt, H. M. (1994). Foreword. In S. Lorenzen (Ed.), *The phylogenetic systematics of free-living nematodes* (pp. xv–xvi). The Ray Society.

Rubin, S., Young, M. H.-Y., Wright, J. C., Whitaker, D. L., & Ahn, A. N. (2016). Exceptional running and turning performance in a mite. *The Journal of Experimental Biology*, 219(Pt 5), 676–685.

Sabatini, M. A., & Innocenti, G. (2001). Effects of Collembola on plant-pathogenic fungus interactions in simple experimental systems. *Biology and Fertility of Soils*, 33(1), 62–66.

Saporito, R. A., Donnelly, M. A., Norton, R. A., Garraffo, H. M., Spande, T. F., & Daly, J. W. (2007). Oribatid mites as a major dietary source for alkaloids in poison frogs. *Proceedings of the National Academy of Sciences of the United States of America*, 104(21), 8885–8890.

Schmelzle, S., & Blüthgen, N. (2019). Under pressure: Force resistance measurements in box mites (Actinotrichida, Oribatida). *Frontiers in Zoology*, 16(1), 24.

Shatilovich, A., Gade, V. R., Pippel, M., Hoffmeyer, T. T., Tchesunov, A. V., Stevens, L., Winkler, S., Hughes, G. M., Traikov, S., Hiller, M., Rivkina, E., Schiffer, P. H., Myers, E. W., & Kurzchalia, T. V. (2023). A novel nematode species from the Siberian permafrost shares adaptive mechanisms for cryptobiotic survival with *C. elegans* dauer larva. *PLoS Genetics*, 19(7), e1010798.

Steel, H., & Ferris, H. (2016). Soil nematode assemblages indicate the potential for biological regulation of pest species. *Acta Oecologica*, 73, 16–24. <https://doi.org/10.1016/j.actao.2016.03.004>

Wu, G. C., & Wright, J. C. (2015). Exceptional thermal tolerance and water resistance in the mite *Paratarsotomus macropalpis* (Erythracaridae) challenge prevailing explanations of physiological limits. *Journal of Insect Physiology*, 82, 1–7.